

### Cervical Cancer Deaths by Race/Ethnicity and Age

Race/ Ethnicity	1992		1993		1994		1995		1996		1996 Age- Adjusted Rate	Calif. 1992 <sup>1</sup>	U.S. 1995	HP 2000 <sup>1,3</sup>
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate				
Overall <sup>α</sup>	44	3.5	32	2.5	35	2.7	41	3.1	21	1.6	1.4	2.6	2.5	1.3
White	24	2.9	18	2.2	21	2.6	20	2.4	14	1.7	1.2	2.1		
Hispanic	10	3.8	9	3.3	7	2.5	11	3.8	2	-	-	3.9	3.5 <sup>2</sup>	2
Black	3	-	3	-	2	-	6	7.9	0	-	-	3.8	5.2	3
Asian/PI	6	6.0	2	-	4	-	3	-	5	4.5	4.2	2.8		
Native American	1	-	0	-	1	-	1	-	0	-	-			
Other	0	-	0	-	0	-	0	-	0	-	-			
Age Groups														
	No.	Rate	No.	Rate	No.	Rate	No.	Rate	No.	Rate				
<1	0	-	0	-	0	-	0	-	0	-				
1-14	0	-	0	-	0	-	0	-	0	-				
15-24	0	-	1	-	0	-	0	-	0	-				
25-64	25	3.9	24	3.7	26	3.9	27	4.1	14	2.1				
65+	19	11.3	7	4.1	9	5.2	14	8.0	7	3.9				

No rates calculated for fewer than 5 events.

<sup>α</sup> Overall may be greater than the sum of cases in Age Groups because deaths in which age could not be determined are included.

Rates calculated per 100,000 population.

Age adjusted rates calculated using the 1940 standard million population.

<sup>1</sup>Age adjusted.

<sup>2</sup>1994 data.

<sup>3</sup>HP 2000 (16.4) "Reduce deaths from cancer of the uterine cervix to no more than 1.3 per 100,000 women."

Source: California Department of Health Services Center for Health Statistics, Vital Statistics Section, Birth and Death Statistical Master Files

**County of San Diego HHS Child, Youth & Family Health Assessment & Planning**